

Table 1. Demographics (Preoperative information)

	n	(%)
Mean age (66.55)		
< 50	1	(1.0)
50-60	26	(25.7)
60-70	37	(36.6)
70-80	27	(26.7)
> 80	7	(6.9)
Gender		
Male	41	(40.6)
Female	60	(59.4)
BMI		
18,5-25	12	(11.9)
25-30	31	(30.7)
30-35	27	(26.7)
35-40	10	(9.9)
>40	5	(5.0)
Missing values	16	(15.8)
Indication		
Pain only	49	(48.5)
Mechanical symptoms	36	(35.6)
Stiffness and pain	13	(12.9)
Other	2	(2.0)
Pain		
Anterior	73	(72.3)
Other	13	(12.9)
Type of pain		
Anterior knee pain	45	(44.6)
Constantly painful knee	22	(21.8)
Other nonspecific symptoms	6	(6.0)
PF-joint incongruent in mountain view radiographs		
Yes	14	(13.9)
No	86	(85.1)
Missing values	1	(1.0)
Osteoarthritis in the PF-joint		
Yes	10	(9.9)
No	90	(89.1)
Missing values	1	(1.0)

Table 2. Peri- and postoperative results.

	n	(%)
Patellar motion (perioperatively)		
Congruent	89	(88.1)
Congruent but rasps	2	(2.0)
Lateralized	7	(6.9)
Missing	3	(3.0)
Complication		
None	85	(84.2)
A bleeding wound/hematoma	8	(7.9)
Infection	5	(5.0)
Satisfaction		
Satisfied with the outcome	48	(47.5)
Better than before surgery	20	(19.8)
Dissatisfied	31	(30.7)
Outcome not known	2	(2.0)
Postoperative symptoms		
No symptoms	39	(39.0)
Residual pain	40	(40.0)
Mechanical symptoms (may also include pain)	7	(7.0)
Instability of the knee	3	(3.0)
Stiffness (may also include pain)	8	(8.0)
Other residual problems	3	(3.0)
Missing	1	(1.0)
Postoperative change		
No symptoms (pain etc.)	39	(38.6)
Residual symptoms (e.g. pain, stiffness etc.)	61	(60.4)
Missing	1	(1.0)